

The *Pseudomonas aeruginosa* PSL polysaccharide is a social but non-cheatable trait in biofilms

Yasuhiko Irie^{1,2*#}, Aled E. L. Roberts^{1§}, Kasper N. Kragh³, Vernita D. Gordon⁴, Jaime Hutchison⁴, Rosalind J. Allen⁵, Gavin Melaugh⁵, Thomas Bjarnsholt^{3,6}, Stuart A. West⁷, Stephen P. Diggle^{1*}

¹ *School of Life Sciences, Centre for Biomolecular Sciences, University of Nottingham, University Park, Nottingham, United Kingdom*

² *Department of Biology & Biochemistry, University of Bath, Claverton Down, Bath, United Kingdom*

³ *Department of International Health, Immunology, and Microbiology, University of Copenhagen, Copenhagen, Denmark*

⁴ *Center for Nonlinear Dynamics and Department of Physics, The University of Texas at Austin, 2515 Speedway, Stop C1610, Austin, TX 78712-1199, USA*

⁵ *School of Physics and Astronomy, University of Edinburgh, James Clerk Maxwell Building, Peter Guthrie Tait Road, Edinburgh EH9 3FD, United Kingdom*

⁶ *Department of Clinical Microbiology, Rigshospitalet, Copenhagen, Denmark*

⁷ *Department of Zoology, University of Oxford, Oxford OX1 3PS, United Kingdom*

*Corresponding authors: steve.diggle@nottingham.ac.uk, yasuhiko.irie@umu.se

#current address: Department of Molecular Biology, Umeå University, 901 87, Umeå, Sweden

§current address: Cardiff School of Health Sciences, Cardiff Metropolitan University, Llandaff Campus, Western Avenue, Cardiff, CF5 2YB, UK

Running title: Social evolution of the biofilm matrix

Author contributions: YI, AELR, and KNK performed research; YI, AELR, KNK, TB, RA, GM, VDG, JH, SAW, and SPD designed the research and analysed the data; YI, SAW, and SPD wrote the manuscript.

Keywords: biofilm, extracellular polysaccharides, social evolution, sociomicrobiology

Abstract

Extracellular polysaccharides are compounds secreted by microorganisms into the surrounding environment and which are important for surface attachment and maintaining structural integrity within biofilms. They have been suggested to be metabolically costly to produce, and because they are secreted, to act as co-operative shared resources within biofilm communities. These assumptions have, however, not been experimentally well-examined. Here we empirically test the cooperative nature of the PSL polysaccharide, which is crucial for the formation of biofilms in *Pseudomonas aeruginosa*. We show that: (1) PSL provides population level benefits in biofilms, for both growth and antibiotic tolerance; (2) the benefits of PSL production are social and are shared with other cells; (3) the benefits of PSL production appear to be preferentially directed towards cells which produce PSL; (4) cells which do not produce PSL are unable to successfully exploit cells which produce PSL. Taken together, this suggests that PSL is a social but relatively non-exploitable trait, and that growth within biofilms selects for PSL-producing strains, even when multiple strains can interact (low relatedness).

The growth and proliferative success of many bacteria, including human pathogens, depends upon their ability to form biofilms in their respective environmental niches. Biofilms are multicellular three dimensional biomass structures, held together by extracellular matrix molecules that encapsulate cells and cause them to aggregate. These extracellular polysaccharides (EPS) which are secreted by the bacteria, typically function as adhesins that are used to attach to a surface and to maintain the three-dimensional biofilm structure, and sometimes aid in protection against a variety of stresses, including dehydration, antibiotics, and predators ^{1, 2}. The production of EPS represents a problem from an evolutionary perspective ³, because it appears to be a type of co-operative behaviour that can potentially provide a benefit to all cells in the community, and not just to those that produce EPS. Consequently, the question arises: “what prevents the invasion of potential cheats that do not produce EPS?”.

A possible solution to this problem is that EPS production may not be an exploitable co-operative trait. Xavier & Foster showed, in an individual based simulation, that if the production of EPS facilitated growth into areas where nutrient availability was greater, then EPS producing lineages could spatially smother and outcompete non-producers ⁴. In this case, EPS production was costly, but the benefits were preferentially provided to nearby cells, which were likely to be EPS-producing clone mates. Some empirical support for this particular mechanism has been demonstrated in *Vibrio cholerae*, where it has been shown that EPS producing lineages are able to displace non-producers ⁵. In addition, EPS producers in *V. cholerae* are also able to share other secreted public goods with each other, which provides another benefit that is preferentially directed towards other EPS producers ^{6, 7, 8}. However, the generality of these explanations for the evolutionary stability of EPS production remains unclear, and the work on *V. cholerae* represents the only empirical study to measure the social costs and benefits of EPS production. In addition, EPS produced by other bacterial species can vary greatly in both their chemical structure and the biological

roles they play within biofilms. Furthermore, many species produce more than one type of EPS that are sometimes but not necessarily co-regulated. This means that there may be differences in the social nature of different types of EPS, and for different bacterial species.

Pseudomonas aeruginosa is an opportunistic pathogen that causes various biofilm infections such as chronic respiratory infections of cystic fibrosis, keratitis, and chronic wound infections. *P. aeruginosa* is known to produce at least three different types of EPS as major components of its biofilm matrix: alginate, PEL, and PSL polysaccharides^{9, 10}. Alginate production is inversely regulated with PSL^{11, 12}, and is not expressed to high levels in the majority of non-CF isolates^{13, 14}. In contrast, PSL is expressed in most *P. aeruginosa* natural and clinical isolates¹³. PSL is a crucial adhesive scaffolding component of the biofilm matrix, promoting both cell-to-cell interactions and surface attachment^{15, 16, 17}. Here we test the social nature of PSL and find that (1) PSL production is not metabolically costly to *P. aeruginosa* cells; (2) PSL+ strains are significantly fitter than PSL- strains in mixed culture biofilms and PSL- strains cannot act as social cheats; (3) the benefit of producing PSL is enhanced when there are many PSL- cells present; (4) biofilms containing a high proportion of PSL- cells are more susceptible to antibiotics; (5) relatedness in biofilms does not matter since PSL+ strains are favoured in conditions of high and low relatedness. More generally we highlight that not all components of the biofilm matrix should be considered as shared resources.

Results

PSL provides a population level benefit in biofilms. We first tested whether PSL provides fitness benefits to a *P. aeruginosa* population growing in biofilms or planktonic/unattached cultures compared to that of populations of mutants that do not produce PSL. In order to simultaneously monitor both unattached and biofilm populations, we modified a previously described bead method¹⁸. This model allowed us to grow biofilms on 7 mm plastic beads in test tubes, harvest biofilm cells from the beads, and directly aspirate unattached cells from the liquid media (Fig. S1). Unattached cells were likely to include planktonic, unattached cellular aggregates, biofilm defective, and/or dispersed cells which can be transient and inseparable or indistinguishable. In this paper we unify them and refer to this sub-population as 'unattached cells'. We used two strains: a PSL+ strain that constitutively produces PSL, and a PSL- mutant that produces no PSL. PSL expression has been shown to induce the intracellular secondary messenger molecule c-di-GMP, which controls multiple biofilm-associated genes¹⁹. Consequently, to ensure that our results are due to PSL and not c-di-GMP, we constitutively elevated c-di-GMP ($\Delta wspF$ background) in both our strains, but nevertheless ultimately found that our results are c-di-GMP-independent because identical results were recorded using non- $\Delta wspF$ backgrounds (Fig. S2a biofilm: $F_{(1,16)} = 17.54$, $p = 0.0007$, $n = 3$; Fig. S2b unattached: $F_{(1,16)} = 0.67$, $p = 0.4251$, $n = 3$; Fig. S2b biofilm: $t(13) = -3.599$, $p = 0.0032$; Fig. S2b unattached: $t(13) = -1.162$, $p = 0.2662$).

Our results show that PSL provides a population level benefit in biofilms but not in unattached populations (Fig. 1a). Consistent with previous reports^{13, 15, 16, 17, 20}, we found that

over 4 days of growth, PSL mutants formed significantly less biomass and therefore poorer biofilms compared to the corresponding wild type (PSL+) ($F_{(1,15)} = 11.26, p = 0.0043, n = 3$). In contrast, we found no significant differences in final population density between the PSL+ and the PSL- strain in the unattached population of cells ($F_{(1,16)} = 0.67, p = 0.4251, n = 3$).

In order to confirm that our unattached population results were not biased by cells that had detached from biofilms, we also performed an experiment where we added no beads and therefore the analyses were all performed on purely unattached cells. Again we found no significant differences in final population density between PSL+ and PSL- strains (Fig. S3). Since we found that a significant biomass was achieved after 1 day of biofilm growth, and in order to minimise the occurrence of rapid spontaneous mutations that can develop and accumulate in mature biofilms of *P. aeruginosa*²⁰, we carried out all subsequent experiments with 24 hour cultures.

PSL provides social benefits in biofilms. We then asked whether the production of PSL provides social benefits to other cells within a biofilm. We tested this in unattached populations and in biofilms, by growing the PSL- strain with either other PSL- cells or PSL+ cells in a 1:1 ratio. We found that approximately 100-fold more PSL- cells could attach and form biofilms in the presence of PSL+ compared to when PSL- was co-cultured with PSL- (Fig. 1b; $p < 0.0001, n = 4$). In contrast, in unattached populations, we found that the fitness of PSL- cells was not influenced by co-culturing with PSL+ (Fig. 1b; $p > 0.08, n = 4$). This indicates that, in biofilms, the production of PSL provides some benefits to cells that do not produce PSL, but not in unattached populations. Consistent with PSL+ providing a benefit to PSL- cells, we found that when we co-cultured PSL+ and PSL- strains on glass microscope slides (rather than beads), PSL- cells co-aggregated with PSL+ cells and incorporated themselves into the biofilm during the early stages of biofilm formation (Fig. 1c).

PSL mutants do not act as social cheats within biofilms. We next tested whether PSL- strains could act as social cheats²¹, i.e. whether these strains increased in frequency when growing in mixed cultures with the PSL+ strain. We varied the starting ratios of PSL-:PSL+ cells from 0.1 to 10,000, because theory predicts that the fitness of cheats should be frequency-dependent, with cheats being better able to exploit co-operators when the cheats are rarer²². In biofilms, we found that the relative fitness of the PSL- cells was either equal to or lower than that of the PSL+ strain, suggesting that the PSL- cells were not able to outcompete the PSL+ cells (i.e. to cheat). The relative fitness of the PSL- strain was negatively correlated with its starting frequency in the population (Fig. 2a biofilm; $t(13) = -3.242, p = 0.0064$). At high starting frequencies of PSL+, the relative fitness of the PSL- strain was not significantly different from that of the PSL+ strain. As the starting frequency of the PSL- strain was increased, the relative fitness of the PSL- strain became lower than that of the PSL- strain. In contrast, we found that in unattached populations, the relative fitness of the PSL- strain did not differ significantly from the PSL+ strain, irrespective of the starting ratio (Fig. 2a unattached; $t(13) = -0.741, p = 0.4716$).

We also tested whether the presence of the PSL- strain adversely affected biofilm production in mixed cultures. Given that PSL- strains are unable to form biofilms with the same biomass as PSL+ strains (Fig. 1), we expected a high starting frequency of PSL- strains would adversely affect biofilm production. In contrast to this expectation, we found that the final biomass did not significantly change as we varied the starting ratio of PSL-:PSL+ cells from 0.1 to 1000 (Fig. 2b biofilm: $t(13) = -1.769$, $p = 0.1004$; Fig. 2b unattached: $t(13) = 1.881$, $p = 0.0825$). This result indicates the final biomass is determined by the PSL+ cells, which dominate long-time biofilm growth independently of the starting ratio between PSL- and PSL+. To allow us to microscopically visualise growth of the PSL- and PSL+ strain in co-inoculated biofilms, we performed a biofilm flow cell experiment with a mixed PSL+ and PSL- population. Consistent with Fig. 1c, PSL- cells appeared to co-aggregate with PSL+ cells and incorporate themselves into the early-stage biofilms (Fig. 3a and b). However, over time, the PSL+ cells covered and outcompeted the PSL- cells (Fig. 3c and d).

PSL provides population level benefits against antibiotics. PSL is known to play a role in tolerance to antibiotics including aminoglycosides^{20, 23} (Fig. S4). We tested the social consequences of this tolerance by examining the effect of adding gentamicin at a concentration previously known to affect biofilm (100 $\mu\text{g/ml}$)²⁴ to populations that contained a variable ratio of PSL- mutants. In the absence of gentamicin, we found that the frequency of PSL- mutants did not influence the biomass of biofilm, consistent with Fig. 2b (Fig. 4a; $t(13) = -0.732$, $p = 0.4769$). In contrast, when we added gentamicin, the biofilm biomass was negatively correlated with the fraction of PSL- mutants in the starting population (Fig. 4a). When there were more PSL- than PSL+ cells in the starting population, the addition of gentamicin led to a significant drop in viable biofilm biomass. In cases where the starting ratio of PSL-:PSL+ was greater than 10, the addition of gentamicin reduced the population to below our threshold detection level. Examining in more detail the populations which showed partial killing, we found that PSL+ cells had a significantly higher level of survival than the PSL- cells (Fig. 4b; $t(4) = 4.07$, $p < 0.02$, $n = 3$).

Relatedness does not influence selection for PSL. We complemented the above fitness assays with a multi-generational selection experiment. A number of previous studies on microbial social traits such as siderophore production and quorum sensing, have shown that the relative fitness of individuals that do and do not perform social traits depends upon population structure^{25, 26, 27, 28}. Specifically, structured populations, with a relatively high relatedness between cells (low strain diversity), favour genotypes that support co-operative traits. Conversely, populations with relatively low relatedness between cells, (high strain diversity) favour less co-operative cheating genotypes. These results are as predicted by social evolution theory^{29, 30}. In contrast, our observation that PSL- is not able to cheat PSL+ in mixed cultures, suggests that relatedness might not matter for the evolution of PSL in the same way.

We tested this hypothesis with a selection experiment over 5 days, in which we started with a mixed population of PSL+ and PSL- cells, and manipulated both relatedness (low/high) and whether cells were growing in biofilms (on beads) or in cultures with no beads (the entire

population is unattached) (Fig. S5). In each round of growth, we subdivided the population into a number of subpopulations. We varied relatedness by initiating each subpopulation with either a single clone, to give relatively high relatedness, or with multiple clones, to give relatively low relatedness. Consequently, in the high relatedness treatment the PSL+ and PSL- strains become segregated into separate populations, while the low relatedness treatment keeps them mixed together in the same population. We started our experiments with a 1:1 ratio of PSL-:PSL+, and carried out six rounds of growth.

We found that, in contrast to previous selection experiments on microbial social traits, relatedness had no significant ($F_{[1, 20]} = 1.69, p > 0.05, n=6$) influence on the outcome of our selection experiment (Fig. 5). Instead, we found that the relative fitness of the PSL- and PSL+ strains were determined by whether the subpopulations were grown as biofilms or in unattached populations. Specifically, the PSL+ strain was significantly ($F_{[1, 20]} = 542.7, p < 0.0001, n=6$) favoured in biofilms but not in unattached populations in bead-less cultures (Fig. 5). Consequently, our selection experiment provides further support for our conclusion that PSL+ is favoured in biofilms, and that PSL- strains are unable to cheat PSL+ strains.

Discussion

EPS are important components of the bacterial extracellular matrix, and EPS production has long been considered to constitute a co-operative social behaviour. Despite this, little empirical work has been performed to test this. We examined the social nature of the PSL polysaccharide produced by *P. aeruginosa* and found that: (1) PSL production provides a benefit at the group level to cells growing in biofilms (Fig. 1a); (2) PSL production provides no benefit or cost to cells growing in unattached populations (Fig. 1a and 2a); (3) PSL- strains are able to grow better as biofilms when in a mixed culture with PSL+ cells (Fig. 1b); (4) PSL+ strains are able to outcompete PSL- strains when in mixed culture biofilms, such that PSL- strains cannot successfully exploit (cheat) the PSL production of the PSL+ strain (Fig. 2a); (5) the relative fitness of the PSL- strain is lower when the PSL+ strain is rarer (negative frequency dependence; Fig. 2a); (6) the presence of PSL- cells has a negligible influence on overall biofilm productivity (Fig. 2b); (7) biofilms containing a higher proportion of PSL+ cells are less susceptible to antibiotics (Fig. 4a); (8) PSL+ cells are better able to survive antibiotics when growing in a mixed culture biofilm with PSL- cells (Fig. 4b); (9) in a selection experiment, the relative success of PSL+ versus PSL- was not influenced by relatedness, for growth in biofilms (Fig. 5).

Overall, our results suggest that in biofilms, the production of PSL is a social trait, which provides benefits at the individual and group level, but that it cannot be successfully cheated by individuals who do not produce it. This is despite the fact that the PSL- strain is able to gain some benefit from the presence of the PSL producing strain in the biofilm matrix (Fig. 1). Our experiments suggest that the fitness benefits of producing PSL accrue mainly to the producing cell and/or to other PSL+ cells. This contrasts with previous studies on social traits in *P. aeruginosa*, where it has been shown that siderophore production and quorum sensing

(QS) are social traits that can be readily exploited by cheats *in vitro*, *in vivo* and in biofilms^{25, 26, 28, 31}. We suggest that this is because siderophores and QS-dependent public goods are relatively freely diffusible secreted products that provide shared benefits to both producing and non-producing cells. In contrast, since its primary function is adhering to cells and attached surfaces, PSL likely has limited diffusive properties in the biofilm biomass. PSL was previously shown to localise primarily to the periphery of the biofilm microcolonies, encapsulating the cells³². In addition, some PSL is known to be tightly associated to the cell surface³³, although some is likely to be diffusible, and serves as an intercellular signalling molecule¹⁹. This signalling property of PSL also promotes co-ordinated, community-level gene expression towards PSL production, since PSL has been shown to promote its own expression in a feed-forward manner, similar to that of an auto-induction regulatory cascade.

Our work shows both similarities and differences with a previous individual-based simulation of biofilms which studied competition between strains that do and do not produce EPS⁴. Specifically, Xavier & Foster⁴: (1) assumed that EPS production is costly, such that where mono-cultures of EPS- strains have a higher growth rate than EPS+; (2) predicted that secretion of EPS allows producing cells to push descendants into more oxygen-rich environments which suffocates non-producers; (3) predicted that EPS producers most easily invade biofilms containing a high proportion of non-producers. We did not find support for Xavier & Foster's assumption that PSL production has a significant effect on bacterial fitness. When we compared PSL+ and PSL- strains in unattached populations, we found no significant differences in growth (Fig. 1a). It is possible that in our culturing condition, growth is limited by other factors, such as the availability of essential minerals and metal ions. In contrast, we did find support for Xavier & Foster's prediction that, in biofilms, PSL producing cells grow over the top of non-producing cells (Fig. 3c and d). However, it is hard to disentangle whether this is a property of the oxygen/nutrient gradient, or whether it is the property of differing adhesive strengths between PSL+ and PSL- cells, as PSL- cells are less adhesive and are more prone to be detached from the biofilm biomass in constant flow conditions when exposed on the surface. A greater adhesive ability can allow strains to displace competitors, and gain an advantage in biofilms⁵.

Our experiments suggest that PSL production cannot be cheated, and hence should be favoured even in low relatedness mixed cultures. Therefore, we tested the effect of relatedness on selection for PSL production in biofilms, using a selection experiment (Fig. 5). In our experiment, high relatedness implied that cultures were established with either PSL+ or PSL- strains. Under these conditions, cells will interact with genetically identical cells, and PSL- cells cannot interact with PSL+ cells. In contrast, low relatedness implies that cultures can be comprised of both PSL+ and PSL- strains, such that PSL- cells can theoretically exploit the PSL+ cells. We found that relatedness had no effect on our results; PSL+ strains were favoured in conditions of high and low relatedness in biofilms but not in non-biofilm populations. Therefore in mixed culture biofilms, a PSL producer always wins, in contrast to previous work on other social traits, such as siderophore production and QS^{25, 26, 28, 31}. In these previous studies, relatedness matters, because at low relatedness, the uncooperative cells are able to interact with the cooperative cells, allowing the uncooperative

1 strains to exploit (cheat) the cooperative wild type cells³⁴. Our selection experiment provides
2 further evidence that the benefits of producing PSL directly affect the producing cells and
3 that PSL production is not exploitable by non-producing cheats.

4
5 We further demonstrated that PSL helps protect biofilm communities against environmental
6 challenge. When biofilms contain a high proportion of PSL⁻ cells, this leads to a significant
7 increase in the susceptibility to antibiotics (Fig. 4a). Within these biofilms, PSL⁺ cells
8 showed a higher level of survival than the PSL⁻ cells (Fig. 4b), indicating that the protection
9 provided to the cells by PSL is most beneficial for the producing cells. This is likely due to
10 PSL being both cell-associated and released into the media¹¹. While PSL⁻ cells can readily
11 access released PSL, they do not have access to cell-associated PSL. One possibility is that
12 there are functional differences for providing antibiotic tolerance between cell-associated
13 PSL and released PSL. Alternatively, the relative PSL concentration may simply be higher
14 in closer proximity to the cell surface, providing a higher level of protection against
15 antibiotics. Shared benefit appears to be provided to the community as a whole including
16 non-producing cells when they are rarer. This may be because biofilm biomass appears to be
17 entirely encapsulated with possibly the released form of PSL³². The exact mechanism of
18 antibiotic tolerance by PSL is not known, and this therefore remains an open question.

19
20 Arguments have been made previously, that *P. aeruginosa* cells, even when they are in the
21 ‘planktonic’ phase, have biofilm-like features^{35,29}. In our bead biofilm model, the unattached
22 cell population, particularly those cells which constitutively express EPS, contain large
23 numbers of cellular aggregates, as seen previously in shaken liquid cultures^{19, 32,18, 26}.
24 However, our results clearly distinguish the social evolutionary differences between the cells
25 in the biofilm and unattached cells, thus providing evidence that unattached cells, whether
26 they are aggregated or not, are definitively distinct from surface-attached biofilm cells. This
27 raises the question whether aggregates represent a type of biofilm, are ‘biofilm-like’, or
28 represent a different growth mode altogether. This is an important question since *P.*
29 *aeruginosa* biofilms growing on infected contact lens, catheters, and other medical implants
30 form surface-associated biofilms, but within the cystic fibrosis sputum, the bacteria are
31 thought to grow as suspended aggregates^{36, 37, 38,30}.

32
33 Most strains of *P. aeruginosa*, like many other microbial species, are capable of producing
34 multiple biofilm EPS molecules^{13, 39,12}. Of these, PSL and PEL are co-regulated by several
35 intracellular regulatory factors including c-di-GMP^{40,31} and both contribute to biofilm
36 development in a majority of *P. aeruginosa* strains including PAO1¹³. Intriguingly, in
37 additional experiments, we found that the social aspects of PEL production are different from
38 those of PSL, pointing towards PEL as being a completely private and non-social good (Fig.
39 S6). This is in contrast to recent work inspecting the competitive fitness of PEL producers
40 using the PA14 strain, which showed community-level benefits in some circumstances⁴¹.
41 PA14 has a deletion mutation in the *psl* promoter through *pslD* gene, and the nature of biofilm
42 formation in this strain, which uses PEL polysaccharide as the sole EPS, appears to have
43 diverged from that of other *P. aeruginosa* isolates. It remains to be seen to what degree in
44 PA14, PEL production has compensated for the loss of PSL on the social evolutionary scale.

These results highlight the importance of recognising diverse roles and effects of different EPS as social traits within biofilms. The extracellular matrix is the material that keeps microbial cells together in a biofilm community, influencing their social interactions and evolution. As such we believe it holds an important foundation for our understanding of the multicellularity of unicellular organisms.

Methods

Bacterial strains and growth conditions. The bacterial strains and plasmids we used and constructed for this study are listed in Table S1. We propagated *Escherichia coli* and *P. aeruginosa* strains in Lysogeny Broth (LB) at 37°C unless otherwise specified. The concentrations of antibiotics we used for *E. coli* were: 50 µg/ml carbenicillin and 10 µg/ml gentamicin, and for *P. aeruginosa*: 300 µg/ml carbenicillin and 100 µg/ml gentamicin. We induced P_{BAD}-*psl* strains for PSL over-expression by adding 1 % w/v L-arabinose. PSL is involved in a feed-forward regulation system where the expression of PSL induces the intracellular levels of c-di-GMP, which in turns over-expresses PSL¹⁹. To avoid c-di-GMP being inadvertently triggered in our biofilm system, which would potentially affect the interpretations of our results when other c-di-GMP-induced gene products become involved, all strains used in our studies were conducted using a Δ *wspF* mutation background. Δ *wspF* mutants express biologically maximal intracellular c-di-GMP levels due to the constitutive activation of diguanylate cyclase WspR⁴².

Fluorescent strain construction. We chromosomally labelled *P. aeruginosa* strains with either GFP or mCherry using the Tn7 delivery system⁴³. We electroporated 1 µl pTNS3 + 1 µl pUC18-mini-Tn7T2-PA1/04/03::*gfp* or pUC18-mini-Tn7T2-PA1/04/03::*mcherry* into a respective electrocompetent *P. aeruginosa* strain as previously described⁴⁴ and selected for gentamicin resistance. We removed the construct backbone by electroporating 1 µl pFLP2 and selecting for carbenicillin resistance. Upon sucrose counter-selection to identify the loss of the *sacB* gene-containing pFLP2 by streaking for single colonies on LB (no salt) plus 10 % sucrose at 30°C, we confirmed the strains for carbenicillin and gentamicin sensitivities.

Bead biofilm culture system. We grew biofilm cultures on plastic beads as previously described¹⁸ but with several modifications (Fig. S1). We used this system because it allowed us to simultaneously study biofilm and unattached populations. This system differs from traditional flow cell models in that it is closed, and unattached populations are not lost due to flow. Therefore any differences that we observe are reflective of biofilm vs unattached populations instead of ‘biofilm cells being the only ones left’, which can occur in a flow cell system. We streaked *P. aeruginosa* strains to obtain single colonies on M9 + 3.062 g/l sodium citrate (M9 citrate) agar at 37°C for 2 days. We picked individual fresh colonies and grew these to logarithmic phase (OD₆₀₀ ≈ 0.5) in M9 citrate broth at 37°C with shaking. We then diluted these cultures to OD₆₀₀ ≈ 0.05 into 3 ml M9 citrate containing one or more (up to three) 7mm plastic beads (Lascells) in standard culture tubes into independent triplicates. We then grew the cultures at 37°C with shaking at 200 rpm for 24 hours. We directly aspirated

unattached cells from the liquid broth portion of the culture. We collected biofilm cells by retrieving the beads from the tubes, and gently washed the beads in 10 ml PBS by five inversions which we repeated three times. We determined the colony forming units (CFU) of the biofilms by recovering the cells by water bath sonication in 1 ml PBS for 10 min. We vortexed the sonicated samples for 10 sec prior to serial dilution and plating.

Antibiotic killing assays. For antibiotic killing assays (Fig. 4), we first grew bead biofilm cultures for 24 hours in the absence of gentamicin. Half the cultures were then introduced to 100 µg/ml gentamicin (final concentration), and the other half were left untreated. We then further incubated the cultures for 24 more hours at 37°C with shaking prior to harvesting the cells for analyses.

Determining ratios of PSL+ and PSL- cells in mixed cultures. To distinguish between PSL+ and PSL- cells, we introduced a single chromosomal copy of constitutively expressed GFP or mCherry genes to each strain. We then used quantitative real-time PCR and primers specifically designed against GFP or mCherry genes to accurately assess the numbers of each strains present in a sample. In order to avoid the effects of PEL, which is co-regulated with PSL, interfering with or convoluting PSL-dependent phenotypes, we conducted all experiments using *Δpel* mutant backgrounds.

Quantitative real-time PCR. We collected genomic DNA (gDNA) from biofilm cells from washed beads without the sonication steps by directly applying lysing reagents to the beads. We isolated unattached bacterial gDNA by pelleting 1 ml of the liquid cultures. We then stored pellets and beads at -20°C until DNA extraction. We extracted gDNA using a GenElute™ Bacterial Genomic DNA Kit (Sigma) according to the manufacturer's protocol except for the elution buffer which we diluted 100-fold to prevent EDTA interfering with the subsequent DNA polymerase reactions. We performed real-time PCR as previously described¹⁹ using Syber Green PCR Master Mix (Applied Biosystems). The primers we used are listed in Table S2. We generated standard curves for *gfp* and *mcherry* genes using quantitatively determined gDNA from *Δpel Δpsl* GFP strain and *Δpel Δpsl* mCherry strain respectively.

Biofilms grown on coverslips and confocal microscopy of biofilms. Bacteria were cultured as described in the bead biofilm system, but instead of inoculating into bead-containing tubes, cultures were diluted into 1 ml M9 citrate in uncoated polystyrene 24-well plates. Each wells contained one sterile 10 mm diameter glass coverslip. Biofilms were grown in 37°C with 200 rpm shaking for 24 hours. The glass coverslips were removed by forceps, gently washed three times in 10 ml PBS, and immediately mounted on a glass slide for confocal microscopy by Zeiss LSM 700. Images were acquired and analysed using Zeiss's ZEN software.

Flow cell growth and confocal microscopy of biofilms. We grew mixed *Δpel Δpsl* and *Δpel* aggregates in a continuous flow cell system as described previously⁴⁵. We produced the mixed aggregates by mixing exponential growing liquid cultures, adjusted to an OD₆₀₀ of 0.001, 1:1 in LB media. This mix was then incubated overnight shaken at 180 rpm. We

diluted the aggregate-containing culture to an OD₆₀₀ of 0.01 before inoculating the flow cells with 27G syringes. We used a flow rate of 3 ml/hour of M9 minimal media buffered with 10 % v/v A10 phosphate buffer supplemented with 0.3 mM glucose. With a Zeiss Imager.Z2 microscope with LSM 710 CLSM and the accompanying software Zeiss Zen 2010 v. 6.0, we found aggregates containing a mix of the GFP- and the mCherry-tagged cells. We imaged these aggregates over time with z-stack intervals of 1 µm. 3D projections of images were produced in Imaris (Bitplane, Switzerland). We quantified biomass of the two tagged populations with the open source software FIJI (LOCI) and the plugin Voxel_Counter (NIH).

Biofilm selection experiment. In order to determine potential fitness of PSL production, we performed a selection experiment over 6 rounds of growth, with four different treatment regimens; relatively low and relatively high relatedness (Lr and Hr, respectively) (Fig. S5). The Hr treatment started with three 3 ml subpopulations of diluted logarithmic phase PSL+ cells and PSL- cells (n = 6). In contrast, the Lr treatment started with six subpopulations of a 3 ml 1:1 mix of PSL+ and PSL- cells (n = 6). We grew cultures at 37°C with shaking at 200 rpm for 24 hours in the presence/absence of plastic beads. We recovered biofilm cells as detailed previously and pooled together prior to serial dilution and plating. We recovered 1ml of unattached cells from the liquid media *in lieu* of plastic beads, and subjected to the same treatment (omitting the three wash stages). We determined the ratio of PSL+ and PSL- cells in each treatment regimen phenotypically; PSL+ and PSL- cells have rugose and smooth colony morphologies, respectively. For Hr treatments we then randomly selected six clones to inoculate 3 ml M9 citrate and progress to the next round, whereas for LR treatments, we randomly took a sweep of multiple colonies. We grew cultures at 37°C with shaking at 200 rpm for 24 hours in the presence/absence of plastic beads adding an additional round of selection. We repeated this experiment six times, with six selection lines per treatment regimen.

Mathematical formula and statistics. We determined relative fitness (w) of the PSL- strain using the following formula: $w = x_2 (1 - x_1) / x_1 (1 - x_2)$, where x_1 is the starting mutant proportion of the population and x_2 is the end mutant proportion²². All error bars denote standard errors of the mean. All statistical tests including unpaired t tests, linear regression analyses and one-way and two-way analyses of variants were performed on VassarStats (<http://vassarstats.net/>).

Acknowledgements

The authors would like to thank Boo Shan Tseng and Matt Parsek for their generous gifts of GFP and mCherry chromosomal labelling constructs. YI would like to thank Dr Andrew Preston for the MC sponsorship and scientific support. This work was supported by a Human Frontier Science Program Young Investigators grant to VG, RJA, TB, and SPD (RGY0081/2012), a NERC grant to SPD and SAW (NE/J007064/1), a European Research Grant to SAW, an FP7 Marie Curie Fellowship grant to YI (PIIF-GA-2012-329832), a grant

from the Lundbeck Foundation to TB and KNK, and Royal Society University Research Fellowships to SPD and RJA.

Competing financial interests

The authors declare no competing or conflicts of financial interests associated with this manuscript.

References

1. Flemming HC. Biofilms and Environmental-Protection. *Water Sci Technol* **27**, 1-10 (1993).
2. Flemming HC, Neu TR, Wozniak DJ. The EPS matrix: the "house of biofilm cells". *J Bacteriol* **189**, 7945-7947 (2007).
3. Boyle KE, Heilmann S, van Ditmarsch D, Xavier JB. Exploiting social evolution in biofilms. *Curr Opin Microbiol* **16**, 207-212 (2013).
4. Xavier JB, Foster KR. Cooperation and conflict in microbial biofilms. *Proc Natl Acad Sci U S A* **104**, 876-881 (2007).
5. Schluter J, Nadell CD, Bassler BL, Foster KR. Adhesion as a weapon in microbial competition. *ISME J* **9**, 139-149 (2015).
6. Drescher K, Nadell CD, Stone HA, Wingreen NS, Bassler BL. Solutions to the public goods dilemma in bacterial biofilms. *Curr Biol* **24**, 50-55 (2014).
7. Nadell CD, Bassler BL. A fitness trade-off between local competition and dispersal in *Vibrio cholerae* biofilms. *Proc Natl Acad Sci U S A* **108**, 14181-14185 (2011).
8. Nadell CD, Drescher K, Wingreen NS, Bassler BL. Extracellular matrix structure governs invasion resistance in bacterial biofilms. *ISME J* **9**, 1700-1709 (2015).
9. Franklin MJ, Nivens DE, Weadge JT, Howell PL. Biosynthesis of the *Pseudomonas aeruginosa* Extracellular Polysaccharides, Alginate, Pel, and Psl. *Front Microbiol* **2**, 167 (2011).
10. Ryder C, Byrd M, Wozniak DJ. Role of polysaccharides in *Pseudomonas aeruginosa* biofilm development. *Curr Opin Microbiol* **10**, 644-648 (2007).

11. Byrd MS, *et al.* Genetic and biochemical analyses of the *Pseudomonas aeruginosa* Psl exopolysaccharide reveal overlapping roles for polysaccharide synthesis enzymes in Psl and LPS production. *Mol Microbiol* **73**, 622-638 (2009).
12. Jones CJ, Ryder CR, Mann EE, Wozniak DJ. AmrZ modulates *Pseudomonas aeruginosa* biofilm architecture by directly repressing transcription of the *psl* operon. *J Bacteriol* **195**, 1637-1644 (2013).
13. Colvin KM, *et al.* The Pel and Psl polysaccharides provide *Pseudomonas aeruginosa* structural redundancy within the biofilm matrix. *Environ Microbiol* **14**, 1913-1928 (2012).
14. Wozniak DJ, *et al.* Alginate is not a significant component of the extracellular polysaccharide matrix of PA14 and PAO1 *Pseudomonas aeruginosa* biofilms. *Proc Natl Acad Sci U S A* **100**, 7907-7912 (2003).
15. Friedman L, Kolter R. Two genetic loci produce distinct carbohydrate-rich structural components of the *Pseudomonas aeruginosa* biofilm matrix. *J Bacteriol* **186**, 4457-4465 (2004).
16. Jackson KD, Starkey M, Kremer S, Parsek MR, Wozniak DJ. Identification of *psl*, a locus encoding a potential exopolysaccharide that is essential for *Pseudomonas aeruginosa* PAO1 biofilm formation. *J Bacteriol* **186**, 4466-4475 (2004).
17. Matsukawa M, Greenberg EP. Putative exopolysaccharide synthesis genes influence *Pseudomonas aeruginosa* biofilm development. *J Bacteriol* **186**, 4449-4456 (2004).
18. Poltak SR, Cooper VS. Ecological succession in long-term experimentally evolved biofilms produces synergistic communities. *ISME J* **5**, 369-378 (2011).
19. Irie Y, *et al.* Self-produced exopolysaccharide is a signal that stimulates biofilm formation in *Pseudomonas aeruginosa*. *Proc Natl Acad Sci U S A* **109**, 20632-20636 (2012).
20. Kirisits MJ, Prost L, Starkey M, Parsek MR. Characterization of colony morphology variants isolated from *Pseudomonas aeruginosa* biofilms. *Appl Environ Microbiol* **71**, 4809-4821 (2005).
21. Ghoul M, Griffin AS, West SA. Toward an evolutionary definition of cheating. *Evolution* **68**, 318-331 (2014).
22. Ross-Gillespie A, Gardner A, West SA, Griffin AS. Frequency dependence and cooperation: theory and a test with bacteria. *Am Nat* **170**, 331-342 (2007).

23. Billings N, *et al.* The Extracellular Matrix Component Psl Provides Fast-Acting Antibiotic Defense in *Pseudomonas aeruginosa* Biofilms. *PLoS Pathog* **9**, e1003526 (2013).
24. Bowler LL, Zhanel GG, Ball TB, Saward LL. Mature *Pseudomonas aeruginosa* biofilms prevail compared to young biofilms in the presence of ceftazidime. *Antimicrob Agents Chemother* **56**, 4976-4979 (2012).
25. Diggle SP, Griffin AS, Campbell GS, West SA. Cooperation and conflict in quorum-sensing bacterial populations. *Nature* **450**, 411-414 (2007).
26. Griffin AS, West SA, Buckling A. Cooperation and competition in pathogenic bacteria. *Nature* **430**, 1024-1027 (2004).
27. Maclean RC, Brandon C. Stable public goods cooperation and dynamic social interactions in yeast. *J Evol Biol* **21**, 1836-1843 (2008).
28. Rumbaugh KP, Trivedi U, Watters C, Burton-Chellew MN, Diggle SP, West SA. Kin selection, quorum sensing and virulence in pathogenic bacteria. *Proc Biol Sci* **279**, 3584-3588 (2012).
29. Hamilton WD. The genetical evolution of social behaviour. I. *J Theor Biol* **7**, 1-16 (1964).
30. Hamilton WD. The genetical evolution of social behaviour. II. *J Theor Biol* **7**, 17-52 (1964).
31. Popat R, Crusz SA, Messina M, Williams P, West SA, Diggle SP. Quorum-sensing and cheating in bacterial biofilms. *Proc Biol Sci* **279**, 4765-4771 (2012).
32. Borlee BR, Goldman AD, Murakami K, Samudrala R, Wozniak DJ, Parsek MR. *Pseudomonas aeruginosa* uses a cyclic-di-GMP-regulated adhesin to reinforce the biofilm extracellular matrix. *Mol Microbiol* **75**, 827-842 (2010).
33. Ma L, Jackson KD, Landry RM, Parsek MR, Wozniak DJ. Analysis of *Pseudomonas aeruginosa* conditional Psl variants reveals roles for the Psl polysaccharide in adhesion and maintaining biofilm structure postattachment. *J Bacteriol* **188**, 8213-8221 (2006).
34. West SA, Griffin AS, Gardner A, Diggle SP. Social evolution theory for microorganisms. *Nat Rev Microbiol* **4**, 597-607 (2006).

35. Schleheck D, *et al.* *Pseudomonas aeruginosa* PAO1 preferentially grows as aggregates in liquid batch cultures and disperses upon starvation. *PloS one* **4**, e5513 (2009).
36. Kragh KN, *et al.* Role of Multicellular Aggregates in Biofilm Formation. *mBio* **7**, (2016).
37. Melaugh G, *et al.* Shaping the Growth Behaviour of Biofilms Initiated from Bacterial Aggregates. *PloS one* **11**, e0149683 (2016).
38. Staudinger BJ, *et al.* Conditions associated with the cystic fibrosis defect promote chronic *Pseudomonas aeruginosa* infection. *Am J Respir Crit Care Med* **189**, 812-824 (2014).
39. Colvin KM, *et al.* The *pel* polysaccharide can serve a structural and protective role in the biofilm matrix of *Pseudomonas aeruginosa*. *PLoS Pathog* **7**, e1001264 (2011).
40. Hickman JW, Harwood CS. Identification of FleQ from *Pseudomonas aeruginosa* as a c-di-GMP-responsive transcription factor. *Mol Microbiol* **69**, 376-389 (2008).
41. Madsen JS, *et al.* Facultative control of matrix production optimizes competitive fitness in *Pseudomonas aeruginosa* PA14 biofilm models. *Appl Environ Microbiol* **81**, 8414-8426 (2015).
42. Hickman JW, Tifrea DF, Harwood CS. A chemosensory system that regulates biofilm formation through modulation of cyclic diguanylate levels. *Proc Natl Acad Sci U S A* **102**, 14422-14427 (2005).
43. Choi KH, *et al.* A Tn7-based broad-range bacterial cloning and expression system. *Nat Methods* **2**, 443-448 (2005).
44. Choi KH, Kumar A, Schweizer HP. A 10-min method for preparation of highly electrocompetent *Pseudomonas aeruginosa* cells: application for DNA fragment transfer between chromosomes and plasmid transformation. *J Microbiol Methods* **64**, 391-397 (2006).
45. Crusz SA, *et al.* Bursting the bubble on bacterial biofilms: a flow cell methodology. *Biofouling* **28**, 835-842 (2012).




a**b****c**

Fig. 1. PSL production provides community benefits to cells grown in biofilms. (a) PSL- strain is significantly defective in biofilm formation on beads compared to PSL+, but no major differences are seen between their growth in the unattached fractions. (b) PSL- co-cultured with PSL+ at a 1:1 ratio increases their proportional numbers in a biofilm (* $p < 0.0001$; ** $p > 0.08$). (c) Confocal micrograph image of surface attached populations of PSL-/PSL+ 1:1 co-cultures. PSL- cells (red) co-aggregate with PSL+ cells (green).



a**b**

Fig. 2. PSL mutants are not social cheats within biofilms. (a) The relative fitness of PSL- cells was equal or lower than the PSL+ strain across all starting frequencies of the mutant in mixed cultures in biofilms. PSL- strain fitness was equal to PSL+ strain fitness across all starting frequencies in unattached populations. (b) PSL-/PSL+ co-culture biomass remained similar across all starting ratios in both biofilm and unattached populations.




Fig. 3. The PSL+ strain smothers and outcompetes the PSL- strain during biofilm growth. Confocal micrograph image of PSL-/PSL+ (1:1) mixed population biofilms at 0 hours (**a** and **b**) and 48 hours (**c** and **d**). PSL- cells are only attached to the surface as co-aggregates with PSL+, and they are eventually outcompeted by PSL+ when the biofilm matures. **a** and **c** are 3D rendered images and **b** and **d** represent corresponding open-box top-down views. Red = PSL+ cells. Green = PSL- cells.



a**b**

Fig. 4. PSL production results in increased tolerance towards antibiotics. (a) Whilst PSL-/PSL+ mixed biofilms yielded consistent biomass regardless of the starting ratio of the two strains, treatment of the mixed biofilm with a high concentration of gentamicin eliminated the entire biofilm population when the starting proportion of PSL- cells increased above PSL+. (b) In the intermediate PSL-:PSL+ co-culture ratio, where mixed biofilm was partially eradicated, there were less PSL- cells in the surviving biofilm in the presence of gentamicin. ($p < 0.02$).




Fig. 5. PSL production is favoured in high and low relatedness conditions in biofilms. PSL+ cells have a selective advantage under conditions promoting biofilm formation (black), however this observation is not reciprocated for unattached conditions (grey). The concurrent loss of PSL- cells from both low related and high related biofilm populations through subsequent selection rounds, suggests there is a strong selection towards PSL+ strains (under biofilm-promoting conditions only) as PSL- strains are unable to cheat on PSL+ strains.

Supplementary Information

Table S1. Bacterial strains and plasmids used in this study.

Strains/plasmids	Description	Source/reference
<i>Pseudomonas aeruginosa</i>		
PAO1	wild type sequenced strain	1
Δpel	defective for PEL production	2
Δpel GFP	Δpel expressing chromosomally labelled constitutive GFP	This study
Δpel mCherry	Δpel expressing chromosomally labelled constitutive mCherry	This study
Δpsl	defective for PSL production	3
Δpsl GFP	Δpsl expressing chromosomally labelled constitutive GFP	This study
Δpsl mCherry	Δpsl expressing chromosomally labelled constitutive mCherry	This study
$\Delta wspF \Delta pel$	constitutively elevated intracellular c-di-GMP; PEL defective	4
$\Delta wspF \Delta pel$ GFP	$\Delta wspF \Delta pel$ expressing chromosomally labelled constitutive GFP	This study
$\Delta wspF \Delta pel$ mCherry	$\Delta wspF \Delta pel$ expressing chromosomally labelled constitutive mCherry	This study
$\Delta wspF \Delta pel \Delta psl$	constitutively elevated intracellular c-di-GMP; defective in both PEL and PSL	2

<i>ΔwspF Δpel Δpsl</i> GFP	<i>ΔwspF Δpel Δpsl</i> expressing chromosomally labelled constitutive GFP	This study
<i>ΔwspF Δpel Δpsl</i> mCherry	<i>ΔwspF Δpel Δpsl</i> expressing chromosomally labelled constitutive mCherry	This study
<i>ΔwspF Δpsl</i>	constitutively elevated intracellular c-di-GMP; PSL defective	4
<i>ΔwspF Δpsl</i> GFP	<i>ΔwspF Δpsl</i> expressing chromosomally labelled constitutive GFP	This study
<i>ΔwspF Δpsl</i> mCherry	<i>ΔwspF Δpsl</i> expressing chromosomally labelled constitutive mCherry	This study
<i>Δpel</i> P _{BAD} - <i>psl</i>	Arabinose-inducible PSL over-expressing strain (<i>Δpel</i> background)	5
<i>Δpel</i> P _{BAD} - <i>psl</i> GFP	<i>Δpel</i> P _{BAD} - <i>psl</i> expressing chromosomally labelled constitutive GFP	This study
<i>Δpel</i> P _{BAD} - <i>psl</i> mCherry	<i>Δpel</i> P _{BAD} - <i>psl</i> expressing chromosomally labelled constitutive mCherry	This study
<i>Δpel Δpsl</i>	defective in both PEL and PSL	2
<i>Δpel Δpsl</i> GFP	<i>Δpel Δpsl</i> expressing chromosomally labelled constitutive GFP	This study
<i>Δpel Δpsl</i> mCherry	<i>Δpel Δpsl</i> expressing chromosomally labelled constitutive mCherry	This study

<i>Escherichia coli</i>		
DH5α	cloning strain F'/endA1 hsdR17(r _k ⁻ m _k ⁺) glnV44 thi-1 recA1 gyrA (Nal ^R) relA1 Δ(lacIZYA- argF)U169 deoR (φ80dlacΔ(lacZ)M15)	Gibco/BRL
XL-10 Gold	cloning strain endA1 glnV44 recA1 thi-1 gyrA96 relA1 lac Hte Δ(mcrCB-hsdSMR-mrr)173 tet ^R F'[proAB lacI ^q ZΔM15 Tn10(Tet ^R Amy Tn5(Kan ^R))]	Stratagene
Plasmids		
pFLP2	FLP recombinase expressing plasmid Amp/Carb ^R	6
pTNS3	Tn7 transposase helper plasmid	7
pUC18-mini-Tn7T2-PA1/04/03::gfp	Suicide vector carrying Tn7 transposable constitutive GFP construct	This study
pUC18-mini-Tn7T2- PA1/04/03::mcherry	Suicide vector carrying Tn7 transposable constitutive mCherry construct	This study

Table S2. Real-time PCR oligonucleotide primers used in this study.

Oligonucleotide primer	Sequence (5' to 3')
<i>gfp</i> RT for1	CGAAAGATCCCAACGAAAAGAG
<i>gfp</i> RT rev1	TCCCAGCAGCTGTTACAAACTC
<i>mCherry</i> RT for1	TTGGACATCACCTCCCACAA
<i>mCherry</i> RT rev1	TCGGCGCGTTTCGTACTGT




Figure S1. Schematic of the bead biofilm system.




Figure S2. PSL-dependent social traits are independent of c-di-GMP. (a) Arabinose-induced PSL over-expressing strain $\Delta pel P_{BAD-psl}$ consistently produced more biofilm biomass compared to the defective mutant $\Delta pel \Delta psl$ ($F_{(1,16)} = 17.54$, $p = 0.0007$). Conversely, no significant differences between the strains were seen in the unattached fraction of the cultures ($F_{(1,15)} = 11.3$, $p > 0.4$). (b) While the experiments in the main figures were performed using $\Delta wspF$ strain background (constitutively elevated c-di-GMP) to remove c-di-GMP and other c-di-GMP-regulated factors from playing a role in our interpretations, repeating the experiment without mutating $wspF$ yielded identical results for both biofilm populations (left panel) and unattached populations (right panel) with the results shown in Fig. 2A, despite $\Delta pel \Delta psl$ having low intracellular c-di-GMP and $\Delta pel P_{BAD-psl}$ having high intracellular c-di-GMP⁵. The cultures were grown in 1% L-arabinose.




Figure S3. EPS mutant strains show no observable differences in growth over four-days in bead-less culture tubes.




Figure S4. PSL expression protects cells from bactericidal concentration of gentamicin. 100 μg/ml gentamicin was added directly to 24 hours bead biofilm cultures and incubated for an additional 24 hours at 37°C with 200 rpm shaking. Resulting populations from both biofilm and unattached cells were tested for viability by serial dilution and plating. ND = not detected.








Figure S5. Schematics of the relatedness experiment. Tubes (populations) were initially inoculated with either single or multiple colonies (equating to populations that had relatively high relatedness [Hr] and relatively low relatedness [Lr], respectively) consisting of PSL+ cells, PSL- cells, or a 1:1 mixture of each. Populations were grown separately in the presence/absence of plastic beads and pooled together before the ratio of PSL+/- cells were determined. Single (Hr) or multiple (Lr) colonies were selected for progression into the next round of the experiment. Should one sub-population (PSL+ or PSL-) have increased fitness, the ratio and thus their likelihood to progress into subsequent rounds of selection will increase.

a**b****c****d**


e

Figure S6. PEL polysaccharides are private goods. (a) Similar to PSL (Fig. 1a), PEL- strain is significantly defective in biofilm formation compared to PEL+ ($F_{(1,24)} = 27.3, p < 0.0001, n = 4$), but no major differences are seen between their growths in the unattached fractions ($F_{(1,24)} = 46.56, p < 0.0001, n = 4$). (b) Unlike PSL, co-cultures of PEL- strains with PEL+ strains do not increase the amount of PEL- strains in neither biofilm nor unattached populations, signifying the unavailability of PEL polysaccharides to non-producing cells. * and ** $p > 0.02$. (c) Confocal micrograph image of surface attached populations of PEL-/PEL+ 1:1 co-cultures. PEL- cells (red) do not co-aggregate with PEL+ cells (green). (d) There is no frequency-dependent fitness changes for PEL-/PEL+ co-cultured biofilm ($t(13) = -0.802, p = 0.4371$), and the relative fitness is consistently slightly below 1, indicating a steady disadvantage of not expressing PEL in the biofilm. There is a frequency-dependent fitness changes in the unattached population ($t(13) = -1.435, p = 0.175$), perhaps caused by complex regulatory system of PEL and the involvement of quorum sensing⁸. (e) Due to PEL not socially affecting co-culture communities, there is a steady decline of biofilm biomass as PEL+ becomes rarer ($F_{(1,8)} = 1, p = 0.441099, n = 3$), but no change is seen in the maximum cell numbers in the unattached population ($F_{(1,8)} = 1.46, p = 0.296592, n = 3$) regardless of the starting ratios of the strains.

References

1. Holloway BW, Krishnapillai V, Morgan AF. Chromosomal genetics of *Pseudomonas*. *Microbiol Rev* **43**, 73-102 (1979).
2. Borlee BR, Goldman AD, Murakami K, Samudrala R, Wozniak DJ, Parsek MR. *Pseudomonas aeruginosa* uses a cyclic-di-GMP-regulated adhesin to reinforce the biofilm extracellular matrix. *Mol Microbiol* **75**, 827-842 (2010).
3. Kirisits MJ, Prost L, Starkey M, Parsek MR. Characterization of colony morphology variants isolated from *Pseudomonas aeruginosa* biofilms. *Appl Environ Microbiol* **71**, 4809-4821 (2005).
4. Irie Y, Starkey M, Edwards AN, Wozniak DJ, Romeo T, Parsek MR. *Pseudomonas aeruginosa* biofilm matrix polysaccharide Psl is regulated transcriptionally by RpoS and post-transcriptionally by RsmA. *Mol Microbiol* **78**, 158-172 (2010).
5. Irie Y, *et al.* Self-produced exopolysaccharide is a signal that stimulates biofilm formation in *Pseudomonas aeruginosa*. *Proc Natl Acad Sci U S A* **109**, 20632-20636 (2012).
6. Hoang TT, Karkhoff-Schweizer RR, Kutchma AJ, Schweizer HP. A broad-host-range Flp-*FRT* recombination system for site-specific excision of chromosomally-located DNA sequences: application for isolation of unmarked *Pseudomonas aeruginosa* mutants. *Gene* **212**, 77-86 (1998).
7. Choi K-H, *et al.* Genetic tools for select-agent-compliant manipulation of *Burkholderia pseudomallei*. *Appl Environ Microbiol* **74**, 1064-1075 (2008).
8. Sakuragi Y, Kolter R. Quorum-sensing regulation of the biofilm matrix genes (*pel*) of *Pseudomonas aeruginosa*. *J Bacteriol* **189**, 5383-5386 (2007).